CHEMOSPHERE

SUBJECT AND AUTHOR INDEX

Volume 22, 1991

Pergamon Press: Oxford New York
Tokyo Seoul

EXECUTIVE EDITOR

Dr. T. Stephen, *8 Lewis Close, Risinghurst, Headington, Oxford OX3 8JD, U.K.*

POLICY BOARD

Chemistry and Biochemistry

— **Professor O. Hutzinger (Chairman) (Alfreda Hutzinger, Editorial Assistant),** *University of Bayreuth, Chair of Ecological Chemistry and Geochemistry, Postfach 10 12 51, D-8580 Bayreuth, G.F.R.*

Ecotoxicology

— **Prof. Dr. J. P. Giesy,** *Department of Fisheries and Wildlife, Michigan State University, East Lansing, MI 48824-1222, U.S.A.*

— **Dr. W. Klein,** *Fraunhofer-Institut für Umweltchemie und Ökotoxikologie, Grafschaft/Hochsauerland, D-5948 Schmallenberg, G.F.R.*

— **Dr. M. Yasuno,** *National Institute for Environmental Studies, Japan Environment Agency, 16-2 Onogawa, Tsukuba, Ibaraki 305, Japan*

Toxicology, Pharmacokinetics and Epidemiology

— **Professor U. G. Ahlborg,** *Karolinska Institutet, Institute of Environmental Medicine, Unit of Toxicology, Box 60208, S-10401 Stockholm, Sweden*

— **Professor S. Safe,** *Veterinary Physiology and Pharmacology, Texas A and M University, College Station, TX 77843, U.S.A.*

— **Dr. E. Takabatake,** *Setsunan University, Faculty of Pharmaceutical Sciences, 45-1 Nagaotoge-cho, Hirakata, Osaka 573-01, Japan*

Atmospheric Chemistry and Global Change

— **Dr. M. A. K. Khalil,** *Oregon Graduate Institute of Science and Technology, 19600 N.W. Von Neumann Drive, Beaverton, OR 97006-1999, U.S.A.*

PARTICIPATING BOARD

Chemistry and Biochemistry

Prof. Dr. K. Bailschmiter, *Universität Ulm (MNH), Abteilung Analytische Chemie, Oberer Eselberg 0 26, D-7900 Ulm, G.F.R.*

Prof. T. F. Bidleman, *Department of Chemistry and Marine Science Program, University of South Carolina, Columbia, SC 29208, U.S.A.*

Dr. R. E. Clement, *Ontario Ministry of the Environment, P.O. Box 213, Resources Road, Rexdale, Ontario, Canada M9W 2L1*

Dr. D. W. Connell, *Griffith University, School of Australian Environmental Studies, Nathan, Brisbane, Queensland 4111, Australia*

Dr. P. E. Des Rosiers, *Research and Development Office, U.S. Environmental Protection Agency, 401 M-Street SW, Washington, DC 20460, U.S.A.*

Dr. H. S. Fuhr, *Bayer AG-Dormagen, Wersverwaltung GEB G-17, D-4047 Dormagen, G.F.R.*

Dr. W. Giger, *Swiss Federal Institute of Technology, Institute for Water Research and Water Pollution Control (EAWAG), CH-8600 Dubendorf, Switzerland*

Prof. Dr. H. P. Hagenmaier, *Department of Organic Chemistry, University of Tübingen, Auf der Morgenstelle 18, D-7400 Tübingen 1, G.F.R.*

Dr. F. Hileman, *Monsanto BB4M, 700 Chesterfield Village Parkway, St Louis, MO 63198, U.S.A.*

Prof. R. A. Hites, *School of Public and Environmental Affairs, 400E 7th Street, Indiana University, Bloomington, IN 47405, U.S.A.*

Prof. Dr. P. M. Huang, *Saskatchewan Institute of Pedology, University of Saskatchewan, Saskatoon, Canada S7N 0W0*

Mr. D. W. Kuehl, *US Environmental Protection Agency, Duluth, MN 55804, U.S.A.*

Dr. R. C. Lao, *Technology Development and Technical Services Branch, Environment Canada, River Road Environmental Technology Centre, Ottawa, Ontario, Canada K1A 0H3*

Prof. D. Mackay, *Department of Chemical Engineering, University of Toronto, Canada M5S 1A4*

Dr. T. Mill, *Stanford Research International, 333 Ravenswood Avenue, Menlo Park, CA 94025, U.S.A.*

Prof. A. A. Moghissi, *PO Box 7166, Alexandria, VA 22307, U.S.A.*

Prof. Dr. H. Parlar, *Gesamthochschule Kassel-Universität, Fachbereich 19 Biologie/Chemie, Heinrich-Plett-Strasse 40, D-3500 Kassel, G.F.R.*

Prof. C. Rappe, *University of Umeå, Institute of Environmental Chemistry, S-901 87 Umeå, Sweden*

Dr. A. Sabljić, *Institute Ruder Boškoyić, Department of Physical Chemistry, P.O.B. 1016, 41001 Zagreb, Croatia, Yugoslavia*

Prof. Dr. H.-R. Schulten, *Fachhochschule Fresenius, Department of Trace Analysis, Dambachtal 20, D-6200 Wiesbaden, G.F.R.*

Dr. G. Sundström, *Dow Sverige AB, P.O. Box 7038, S-103 86 Stockholm, Sweden*

Prof. Dr. P. R. Wallnöfer, *Bayerische Landesanstalt für Ernährung, Menzinger Strasse 54, D-8000 München, G.F.R.*

Dr. V. Zitko, *Program Head, Toxicology and Organic Contaminants Section, Marine Chemistry Division, Department of Fisheries and Oceans, Biological Station, St. Andrews, New Brunswick, Canada E0G 2X0*

Ecotoxicology

Dr. S. M. Bartell, *Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37830, U.S.A.*

Dr. G. C. Butler, *4694 West 13th Avenue, Vancouver, British Columbia, Canada V6R 2V7*

Prof. Dr. D. Calamari, *Istituto di Entomologia, Facoltà di Agraria, Università degli Studi di Milano, Via Celoria 2, 1–20133 Milano, Italy*

Dr. R. T. Digiulio, *Duke University, School of Forestry and Environmental Studies, Durham, NC 27706, U.S.A.*

Prof. Dr. W. Ernst, *Alfred-Wegener-Institut für Polar- und Meeresforschung, Postfach 120161, Columbusstrasse, D-2850 Bremerhaven, G.F.R.*

Prof. M. Goto, *Institute of Ecotoxicology, Gakushuin University, Faculty of Science, 151 Merjiro, Toshima-Ku, Tokyo 171, Japan*

Dr. P. C. Kearney, *Natural Resources Institute, USDA, ARS, Office of Deputy Area Director, Building 001, Room 233, Barc-West, Beltsville, MD 20705, U.S.A.*

Dr. S. J. Klaine, *Memphis State University, Department of Biology, Memphis, TN 38152, U.S.A.*

Dr. P. F. Landrum, *Great Lakes Environmental Research Laboratory, NOAA/USDC, 2205 Commonwealth Boulevard, Ann Arbor, MI 48105–1593, U.S.A.*

Dr. R. Nagel, *Johanne Gutenberg-Universität Mainz, Fachbereich Biologie, Institut für Zoologie, Postfach 3980, Saarstrasse 21, D-6500 Mainz, G.F.R.*

Prof. F. Schmidt-Bleek, *OECD 2, Rue André-Pascal, F-75775 Paris Cedex 16, France*

Dr. A. Spacie, *Perdue University, Department of Forestry and Natural Resources, Forestry Building, West Lafayette, IN 47907, U.S.A.*

Prof. R. Truhaut, *Université de Paris, 4 Avenue Del Observatory, Centre Faculté de Pharmacie, F-75006 Paris 6, France*

Toxicology, Pharmacokinetics and Epidemiology

Dr. R. Kociba, *Dow Chemical Company, 1803 Building, Midland, MI 48674, U.S.A.*

Prof. Dr. Y. Masuda, *Daiichi College of Pharmaceutical Sciences, 22–1 Tamagawa-Cho, Minami-ku, Fukuoka 815, Japan*

Dr. W. Mücke, *Ludwig-Maximilians Universität, Bavarian State Ministry for State Development and Environmental Affairs, Seefeldstrasse 11, D-8018 Grafing, G.F.R.*

Dr. H. Nakazawa, *Laboratory of Medicinal Chemistry, Department of Pharmaceutical Sciences, Institute of Public Health, 6-1, Shirokanedai 4 Chome, Minato-ku, Tokyo, 108 Japan*

Prof. Dr. Ch. Schlatter, *Institute of Toxicology, Federal Institute of Technology, University of Zurich, Schorenstrasse 16, CH–8603 Schwerzenbach, Switzerland*

Prof. Dr. R. R. Suskind, *Department of Environmental Health, Kettering Laboratory, University of Cincinnati, 3223 Eden Avenue, Cincinnati, OH 45267, U.S.A.*

Atmospheric Chemistry and Global Change

Dr. P. Brimblecombe, *School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, U.K.*

Prof. C. I. Davidson, *Department of Civil Engineering, Carnegie Mellon University, Schenley Park, Pittsburgh, PA 15213, U.S.A.*

Dr. J. W. Elkins, *United States Department of Commerce, NOAA–ERL, 325 Broadway, Mail Stop R/E/AR4, Boulder, CO 80303-3328, U.S.A.*

Dr. J. Fishman, *NASA Langley Research Center, Mail Stop 401A, Hampton, VA 23665-5225, U.S.A.*

Dr. S. A. Penkett, *School of Environmental Sciences, University of East Anglia, Norwich NR4 7TJ, U.K.*

Prof. Dr. W. Seiler, *Fraunhofer Institute for Atmospheric Environmental Research, Kreuzeckbahnstrasse 19, D-8100 Garmisch-Partenkirchen, G.F.R.*

Prof. J. W. Winchester, *Department of Oceanography B-169, Florida State University, Tallehassee, FL 32306, U.S.A.*

Twenty-four issues per year, published in two volumes.

Copyright © 1991
Pergamon Press plc

SUBJECT INDEX VOL. 22, 1991

Corrigenda
Addink R., van Bavel B., Visser R., Wever H., Slot P. and Olie K.: 20,1929(1990): conclusions drawn from experiment of VIII with pentachlorobenzene are not supported by experimental evidence. The experiment will be repeated with pure pentachlorobenzene, 799
Trapp,St. and Pussemier L.: 22, 327(1991): on p335–336, correct formula for uptake into stems is given, 1097

Absorption
and catalytic properties, montmorillonite, reaction with organic compds, 769
gravimetric–, and FTIR analysis, interaction and formation of ethoprophos–grafitic oxide interlayer compds, 905

Activated carbon
re–activation by chemchar process, 1191

Adenine nucleotides
detn method of singlet oxygen by use of α–terpinene, 1211

Algae
Selenastrum capricornutum, relationship between bioconcentration and toxicity of volatile aromatic hydrocarbons, 665

Alkenes
reaction with montmorillonite, adsorptive and catalytic properties, 769

Allometry
and individual response, examination of field data for sublethal toxicological effects, 747

Ames test
cyclic and linear tetramines, reversal of nickel mediated biochemical alterations, 272
raw and treated water extracts, genotoxic activity of ether insoluble organic compds, 1089

Amines
ten anilines, uptake, elimination, and bioconcentration in *Brachydanio rerio*, 351

Animals
Baltic, Finnish and Arctic wildlife, chloroterpenes and other organochlorines, 47
dolphins from 1987/1988 mass mortality, US Atlantc coast, presence of chemical residues, 1083
feral mice from PCDD/PCDF contaminated area, induction of hepatic P450–dependent monooxygenase, 1011
herring gull eggs from Great Lakes, re–analysis of archived specimens, comparison of PCB quantitation methods, 201
individual response and allometry, examination of field data for sublethal toxicological effects, 747
male Fischer 344 rats, metabolism of 2,5–dimethylhexane, 77
mallards and bobwhites exposure to compd 1080, acute toxicity and food avoidance behaviour, 655
mammals, predictive QSAR model for evaluating side effects of phenylureas, 613
rats, acidotic, effect of plasma from several tissues on phosphate dependent glutaminase activity, 521
rats, enhancement of fecal excretion of PCBs by addition of rice bran fiber to diet, 375
rats, kerosene treated, enzyme activity in lungs, 705

Analysis
2,3,7,8–TCDF and 2,3,7,8–TCDD, part per quadrillion levels in polyethylene, procedure and initial results, 215
and collection, sorbents for trace levels of airborne bis(2–chloroethyl)sulfide, 461
asbestos in sludges from 51 cities in USA, 963

bootstrap method, aqueous solubility estimated from n–octanol/water partition coeff, 953
chloroaromatic products formed by irradiation of ZnO with aqueous soln of 2,4–dichlorophenol, 1053
coplanar PCB residues in human adipose tissue from Ontario municipalities, 1019
gravimetric adsorption and FTIR analysis, interaction and formation of ethoprophos–grafitic oxide interlayer compds, 905
isotope dilution HRGC/HRMS, coplanar PCBs in fish and marine animal tissue, 849
mass spectrometry and multidimensional chromatography, oil products, pyrolysis of polyethylene–polystyrene mixtures, 887
organochlorines and chloroterpenes in Baltic, Finnish and Arctic wildlife, 47
polycyclic aromatic hydrocarbon content of atmospheric particulate, fuzzified chromatograms for monitoring changes, 1103
quantitative measurement and identification of organic constituents in hazardous waste dumps, 913
radioactively labelled 4,5,6–trichloroguaiacol and fish bile for monitoring aquatic pollutants, 39
re–analysis of archived specimens of herring gull eggs from the Great Lakes, comparison of PCB quantitation methods, 201
sorption of nonpolar and polar compds, applicability of modified OECD–guideline, processes, measurements and experience, 106
speciation and quantitation of Aroclors in hazardous wastes based on PCB congener data, 147
sublethal ecotoxicological effects using allometry and individual biota transformations, 747
toxaphene in fish from Northern Europe, 57

Aquatic organisms see also Fish
fish, crabs and lobsters, PCDD and PCDF contamination, levels and patterns, Newark Bay and New York Bight, 239
marine animal tissue, isotope dilution HRGC/HRMS, coplanar PCBs, 849
monitoring pesticide toxicity from river water, using *Paratya compressa improvia*, 229
mussels, uptake of organochlorines from aquatic ecosystems, 997

Aquatic systems see also Water
aquatic microcosm with different alkalinity and hardness, toxicity and alkalinity of Cu, 577
ecosystems, uptake of organochlorines by hexane–filled dialysis membranes and by mussels, 997
estuarine sediments, butyltins two years after tributyltin use was restricted, 1113
river water, monitoring pesticide toxicity to aquatic organisms using *Paratya compressa improvia*, 229
rivers Rhine and Neckar, PCDDs and PCDFs in fish, 29
sediments in model systems and clay minerals, sorption of HEDP and NTA, 15
soft water, short–term toxicity of fluoride ion to rainbow and brown trout, 603
Thermaikos gulf, Greece, runoff losses of herbicides, 645
USA great lakes, concentration inorganic and organic contaminants. 809

Aquifer material
low organic–carbon–, nonequilibrium sorption of organic chemicals, 341

Asbestos
in sewage sludges from 51 cities in USA, 963

Atmosphere see also Air
biohalogenation as source of halogenated anisoles, 557
deposition of organo chlorine contaminants to Arctic, estimation, 165
particulate, changes in polycyclic aromatic hydrocarbon content monitored by fuzzified chromatograms, 1103
pollution by auxin–type herbicides in Tala valley, Natal, 137
trace levels of airborne bis(2–chloroethyl sulfide, sorbents for analysis and collection, 461
urban–, phase distribution and particle size of polycyclic aromatic hydrocarbons, 503

Bacteria
bacterial mutagen 3–chloro–4–(dichloromethyl)–5–hydroxy–2(5H)–furanone and related compds, formation via chlorination of phenols, 547
liquid growth medium and soil, extraction of pentachlorophenol, comparison of methods used, 99

Behavior
individual response and allometry, examination of field data for sublethal toxicological effects, 747
mallards and bobwhites exposed to compd 1080, acute toxicity and food avoidance, 655

Bioaccumulation
chemical residues, in dolphins from 1987/1988 mass mortality, US Atlantc coast, 1083

Bioassay
polycyclic organic compds in flyash using cytochrome P450IA1 induction, mouse hepatoma cell culture, 895
sediments, effect of storage time on toxicity of *D.magna* and *C.tentans*, 801

Biodegradation
and toxicity of volatile aromatic hydrocarbons, relationship between by *Selenastrum capricornutum*, 665
toxaphene in fish from N Europe, 57
uptake, and elimination of ten anilines by *Brachydanio rerio*, 351

Biodegradation
PCBs (Fenclor 42) in batch cultures with mixed and pure aerobic cultures, 3

Biotransformation
comparative biodegrad–ability, anionic surfactants in synthetic and natural test systems, 873

Biphenyls
aroclor 1254, acetone–induced photode–chlorination in alkaline 2–propanol, mechanism by thermolysis in the presence of di–t–butyl peroxide, 1161
coplanar PCB residues in human adipose tissue from Ontario municipalities, 1019
coplanar PCBs in fish and marine animal tissue, isotope dilution HRGC/HRMS, 849
Fenclor 42 in batch cultures with mixed and pure aerobic cultures, biodegradation, 3
PCB congener data, basis for speciation and quantitation of Aroclors in hazardous wastes, 147
PCB quantitation methods, re–analysis of archived herring gull eggs from Great Lakes, 201
PCB, 'de novo' synthesis in pilot incinerator, 1045
PCBs, aqueous solubility and octanol–water partition coeff, prediction, 723
PCBs, enhancement of fecal excretion by addition of rice bran fiber to diet in rats, 375
PCBs, partition coeff in 1–octanol/water, critical evaluation of measurement, 1175
PCBs, PCFs and PCDs, retention in sediments, motor oil and fats, 421
PCBs, photodechlorination by sodium methyl siliconate, 189

Bleach and bleaching
spent bleach liquor, 2,3,5–trichloro– and 2,3,6–trichloro–p–cymene ne, FTIR and mass spectral data, 881

Carbamates
uptake and translocation of–, by bean plants, model calculations and measurements, 327

Carboxylic acids
4,4'–dichlorobenzilic acid, urinary excreation by dicofol pesticide applicators, 365

Carcinogens
aqueous solubility and octanol–water partition coeff, prediction, 723
ether insoluble organic compds in raw and treated water extracts, 1089

Catalysts
and adsorption of montmorillonite in reaction with organic compds, 769

Chelating agents
HEDP and NTA, sorption onto clay minerals and sediments in aquatic model systems, 15

Chrmatography
2,5–dimethylhexane, metabolism in male Fischer 344 rats, 77
chloroaromatic products formed by irradiation of ZnO with aqueous soln of 2,4–dichlorophenol, 1053
composition of polycyclic aromatic hydrocarbon mixtures in uncontrolled hazardous waste dumps, 317
coplanar PCB residues, human adipose tissue from Ontario municipalities, 1019
FTIR and mass spectral data, 2,3,5–trichloro–and 2,3,6–trichloro–p–cymene ne in spent bleach liquor from activated sludge, 881
fuzzified–, changes in polycyclic aromatic hydrocarbon content of atmospheric particulate, 1103
gas–, retention indices for selected halogenated dioxins, 455
high performance size exclusion–, automated detn of dioxins and furans in fats motor oil and sediments, 437
high performance size exclusion–, effect on retention of dioxins, furans and PCBs in biological samples, 421

isotope dilution HRGC/HRMS, coplanar PCBs in fish and marine animal tissue, 849
multidimensional–, and mass spectrometry, analysis oil products, pyrolysis of poly–ethylene–polystyrene mixtures, 887
quantitative measurement and identification of organic constituents in hazardous waste dumps, 913
speciation and quantitation of Aroclors in hazardous wastes based on PCB congener data, 147
tetracyclic aromatic hydrocarbons, reactivity and product distribution in aqueous chlorination, 495
toxaphene in fish from Northern Europe, 57

Clays
clay minerals and sediments in aquatic model systems, sorption of HEDP and NTA, 15
montmorillonite, reaction with organic compds, adsorptive and catalytic properties, 769

Climate
global consequences of increasing tropospheric ozone concentrations, 685
increasing concentrations produced by human activity, radiative convective climate model, 383
radiative convective model, one dimensional time dependent, theory and development, 383

Clothing
deltamethrin applied to cotton fabric for tsetse fly control, factors affecting its persistence, 677

Combustion
pentachlorophenol containing solids and liquids, formation of PCDDs/PCDFs, 1029
turbulent flame–, of methane with HCl presence of chlorinating species, 67

Concentration
inorganic and organic contaminants. USA great lakes, 809
steady–state–, OH radicals in titanium dioxide aqueous suspension using n–butyl chloride as probe, 1003

Contamination
food–, discrimination by mallards and bobwhites exposed to compd 1080, 655

Cultures
batch cultures of PCBs biodegradation with mixed and pure aerobic cultures, 3
mouse hepatoma cells, bioassay of polycyclic organic compds in flyash using cytochrome P450IA1, 895

Cuticles
of fruit and leaves. permeation and sorption of phenylurea herbicides, effects of cuticular characteristics and climatic parameters, 85

Cyanides
cyanide ion or cyanogen chloride, formation via cleavage of polyhydroxy benzenes with nitrous acid, 267

Degradation
aqueous nitrophenols and nitrobenzene by Fentons reaction, 529
photo–, brominated dibenzodioxins and furans in organic solvents, 821
tributyltin–, butyltins in estuarine sediments two years after use of TBT was restricted, 1113

Dehalogenation
acetone–induced photodechlorination of aroclor 1254 in alkaline 2–propanol, mechanism by thermolysis in the presence of di–t– butyl peroxide, 1161
dechlorination of chloroaromatic compds, 959
photodechlorination, chlorobiphenyls by sodium methyl siliconate, 189

Deposition
air pollution in Arctic by organochlorine compds, 165

Diet see also Food
addition of rice bran fiber, enhancement of fecal excretion of PCBs by rats, 375

Dioxins
2,3,7,8–TCDD excretion, Vietman veterans model for estimating prior exposure to herbicide, 597
2,3,7,8–TCDD, part per quadrillion levels in polyethylene, procedure and initial results, 215
and furans in fats motor oil and sediments, chromatographic automated detn, 437
brominated dibenzodioxins, photodegradation in organic solvents, 821
bromo– and bromochloro–dioxins, retention indices by gas chromatography, 455
chlorinating species, presence in turbulent flame combustion of methane with HCl, 67
concentration on contaminated surfaces, reduction, 485
PCDD contamination, fish, crabs and lobsters, levels and patterns, Newark Bay and New York Bight, 239

PCDD *de novo* synthesis, effects of Cu, Na, K and Zn in solid waste flyash from incinerators, 473
PCDD emissions from waste incinerator, reduction by full scale denox unit, 569
PCDD, 'de novo' synthesis in pilot incinerator, 1045
PCDD, mechanism of formation from flyash and copper chloride, 987
PCDD contaminated area, induction of hepatic P450–dependent monooxygenase in feral mice, 1011
PCDD, formation by pyrolysis and combustion of pentachlorophenol containing solids and liquids, 1029
PCDD formation from bromine precursors, 1121
PCDD in cooking oil causing Yusho disease, biological effects, 537
PCDD in fish from rivers Rhine and Neckar, 29
PCDD in flyash leached from incinerators to water system, 1071
PCDD, partition coeff in 1–octanol/water, critical evaluation of measurement, 1175
PCDs, PCFs and PCBs, retention in sediments, motor oil and fats. 421

Diseases
Yusho–, biological effects of polychlorinated compds in cooking oil, 537

Distribution
and toxicity of Cu in aquatic microcosm under different alkalinity and hardness, 577

Drugs
cyclic and linear tetramines for chelating drugs, reversal of nickel mediated biochemical alterations, 272

Dyes
water soluble azo dyes, fate in activated sludge process, 107

Ecological studies
examination of toxicological data using allometry and individual biota transformations, 747

Effluents
industrial–, in France, biological aspects of regulations, 625

Elimination
chick–, effects of PCDFs and PCDDs in cooking oil causing Yusho disease, 537
uptake, and bioconcentration of ten anilines by *Brachydanio rerio*, 351

Environment
comparative biodegradability of anionic surfactants, synthetic and natural test systems, 873
factors, formation by human lymphoblastic cells and drug metabolism, relationship with environmental factors and aryl hydrocarbon hydroxylase activity, 739
global consequences of increasing tropospheric ozone concentrations, 685
PCDD/PCDF contaminated area, induction of hepatic P450–dependent monooxygenase in feral mice, 1011
PCDDs/PCDFs in flyash leached from incinerators to water system, 1071
sediment, fish tissue and ground water, STORET data, toxic deposition levels, regional variation, USA, 835
sublethal toxicological effects, analysis using allometry and individual response, 747

Enzymes
activity in lungs, kerosene treated rats, 705
aryl hydrocarbon hydroxylase activity, relationship with drug metabolism and fluorescein formation by human lymphoblastic cells, 739
cytochrome P450IA1, induction of mouse hepatoma cell culture, bioassay of polycyclic organic compds in flyash, 895
hepatic P450–dependentmonooxygenase, induction of feral mice from PCDD/PCDF contaminated area, 1011
phosphate dependent glutaminase activity, effect of plasma from several tissues of acidotic rats, 521

Esters and Esterification
montmorillonite as effective catalyst, 769

Ethers and Etherification
insoluble organic fraction of raw and treated water extracts, genotoxic activity, 1089
montmorillonite as effective catalyst, 769

Excretion
2,3,7,8–TCDD–, Vietman veterans model for estimating prior exposure to herbicide, 597
fecal–, enhancement of PCBs by the addition of rice bran fiber to diet in rats, 375
urinary–, 4,4'–dichlorobenzilic acid by applicators of dicofol pesticide, 365

Exposure
mallards and bobwhite–, to compd 1080, acute toxicity and food avoidance behaviour, 655

Extraction
pentachlorophenol from aqueous and soil systems, comparison of methods used, 99
solid phase–, leachability of polycyclic aromatic hydrocarbons from waste, 713

Fats
chromatographic automated detn of furans and dioxins, 437
retention of PCFs, PCDs and PCBs, 421

Filters
adsorption of polycyclic aromatic hydrocarbons, 713

Fish
bile as a tool for monitoring aquatic pollutants using radioactively labelled 4,5,6–trichloroguaiacol, 39
Brachydanio rerio, uptake, elimination, and bioconcentration of ten anilines,351
crabs and lobsters, PCDD and PCDF contamination, levels and patterns, Newark Bay and New York Bight, 239
fish and marine animal tissue, isotope dilution HRGC/HRMS, coplanar PCBs, 849
from Northern Europe, detn of toxaphene in, 57
from rivers Rhine and Neckar, PCDDs and PCDFs, 29
rainbow and brown trout, short–term toxiccity of fluoride ion in soft water, 603
tissue STORET data, toxic deposition levels, regional variation, USA, 835
toxicity of α–terthienyl in near UV–light in lab decreases with time, 699

Fluorescein
formation by human lymphoblastic cells and drug metabolism, relationship with environmental factors and aryl hydrocarbon hydroxylase activity, 739

Fly ash
aromatic bromine compds adsorbed on–, surface catalyzed halogenation–dehalogenation reactions, 1121
concentration and cumulative emission of polycyclic aromatic hydrocarbons, 713
mechanism of formation of PCDD/PCDF with copper chloride, 987
PCDDs/PCDFs leached from incinerators to water system, 1071
polycyclic organic compds, bioassay using cyto–chrome P450IA1 induction, mouse hepatoma cell culture, 895
solid waste incinerator–, effect of Cu, Na, K and Zn on PCDD formation, 473

Food
avoidance and acute toxicity of compd 1080 to mallards and bobwhites, 655

Fungicides
chloroaromatic products formed by irradiation of ZnO with aqueous soln of 2,4–dichlorophenol, 1053
pentachlorophenol, extraction from aqueous and soil systems, comparison of methods used, 99

Furans
2,3,7,8–TCDF, part per quadrillion levels in polyethylene, procedure and initial results, 215
and dioxins in fats motor oil and sediments, chromatographic automated detn, 437
brominated dibenzofurans, photodegradation in organic solvents, 821
chlorinating species, presence in turbulent flame combustion of methane with HCl, 67
PCDF concentration on contaminated surfaces of PCB capacitator, reduction, 485
PCDF contaminated area, induction of hepatic P450–dependent monooxygenase in feral mice, 1011
PCDF contamination, fish, crabs and lobsters, levels and patterns, Newark Bay and New York Bight, 239
PCDF, 'de novo' synthesis in pilot incinerator, 1045
PCDF emissions from waste incinerator, reduction by full scale denox unit, 569
PCDF, formation by pyrolysis and combustion of pentachlorophenol containing solids and liquids, 1029
PCDF, formation from bromine precursors, 1121
PCDF in cooking oil causing Yusho disease, biological effects, 537
PCDF in fish from rivers Rhine and Neckar, 29
PCDF in flyash leached from incinerators to water system, 1071
PCDF mechanism of formation from flyash and copper chloride, 987
PCFs, PCDs and PCBs, retention in sediments, motor oil and fats, 421

Halogen compounds
aromatic bromine compds adsorbed on fly ash, surface catalyzed halogenation–dehalogenati on reactions, 1121

chlorinated benzenes, partition coeff in 1–octanol/water, critical evaluation of measurement, 1175
chlorinated–benzene sorption to hydrophobic surfaces, temp effects, 1141
chloroaromatic compds, use of visible light in photoreduction, 959
chlorodihydroxybenzene formed by photo–transformation of monochlorophenols induced by excitation of nitrate ions, 1063
halogenated benzenes, aqueous solubility and octanol–water partition coeff, prediction, 723
halogenated hydrocarbons, formation from methane, HCl and oxygen, effect of time, temp and reagent concentration, 929
organochlorines and chloroterpenes in Baltic, Finnish and Arctic wildlife, 47
organochlorines, uptake from lake water by hexane–filled dialysis membranes and by mussels, 997
polybrominated chemical residues, in dolphins from 1987/1988 mass mortality, US Atlantc coast, 1083
polychloroinated aromatic compds, formation with copper chloride, mechanism, 987

Halogenation
aqueous chlorination, tetracyclic aromatic hydrocarbons, reactivity and product distribution, 495
biohalogenation as source of halogenated anisoles in air, 557
chlorinating species, presence in turbulent flame combustion of methane with HCl, 67
chlorination, phenols, formation of 3–chloro–4–(dichloromethyl)–5–hydro xy–2(5H)–fur anone and related compds, 547
halogenation–dehalogenatio n, surface catalyzed reactions of aromatic bromine compds adsorbed on fly ash, 1121

Halogens
fluoride ion in soft water, short–term toxiccity to rainbow and brown trout, 603

Health hazards see also Risk assessment
4,4'–dichlorobenzilic acid, urinary excreation by dicofol pesticide applicators, 365
asbestos in sludges from 51 cities in USA, 963
bis(2–chloroethylsulfide, airborne, sorbents for analysis and collection, 461
dermal transfer of surface pesticide residue from indoor fogger use measured by CDFA method, 975

Herbicides
2,3,7,8–TCDD excretion, Vietman veterans model for estimating prior exposure to herbicide, 597
auxin–type–, pollution in Tala valley, Natal, 137
chloroaromatic products formed by irradiation of ZnO with aqueous soln of 2,4–dichlorophenol, 1053
phenylurea–, permeation and sorption of cuticles of fruit and leaves. effects of cuticular characteristics and climatic parameters, 85
runoff losses into Thermaikos gulf, Greece, 545

Humic substances
humic acid in soil, interaction and formation of ethoprophos–grafitic oxide interlayer compds FTIR and gravimetric adsorption analysis, 905

Humidity
effect on penetration of phenylurea herbicides from a droplet through cuticles of fruit and leaves, 85

Hydrocarbons
2,5–dimethylhexane, metabolism in male Fischer 344 rats, 77
benezene–naphthalene mixtures, pyrolysis, 279
methane, turbulent flame combustion with HCl, presence of chlorinating species, 67
polycyclic aromatic hydrocarbon mixtures, composition of in uncontrolled hazardous waste dumps, 317
polycyclic aromatic hydrocarbon content of atmospheric particulate, fuzzified chromatograms for monitoring changes, 1103
polycyclic aromatic–, aqueous solubility and octanol–water partition coeff, prediction, 723
polycyclic aromatic–, in urban atmosphere, phase distribution and particle size, 503
polycyclic aromatic–, leachability from waste by solid phase extraction, 713
products, pyrolysis of polyethylene–polystyrene mixtures, mass spectrometry and multidimensional chromatography, 887
tetracyclic aromatic–, reactivity and product distribution in aqueous chlorination, 495

volatile aromatic–, bioconcentration and toxicity, relationship between by *Selenastrum capricornutum*, 665

Incinerators
modular–, phenomeno–logical models of chlorinated hydrocarbon emissions, 121
PCDD and PCDF emission levels before and after SCR–DeNox–unit, 569
pilot–, 'de novo' synthesis of PCDD, PCDF, PCB, PCN and PAH, 1045

Industry
effluents in France, biological aspects of regulations, 625

Irradiation
malfunctioning PCB capacitator, concentration of PCDD and PCDF on contaminated surfaces, 485
ZnO, with aqueous soln of 2,4–dichlorophenol, chloroaromatic products analysed by HPLC and GC–MS, 1053

Insecticides
deltamethrin applied to cotton fabric for tsetse fly control, factors affecting its persistence, 677
pentachlorophenol, extraction from aqueous and soil systems, comparison of methods used, 99

Insects
tsetse fly control, deltamethrin applied to cotton fabric, factors affecting its persistence, 677

Isotopic labelling
radioactively labelled 4,5,6–trichloroguaiacol and fish bile for monitoring aquatic pollutants, 39

Kinetics
nonequilibrium sorption, organic chemicals by low organic–carbon aquifer materials, 341

Laws
regulations in France, biological aspects of industrial effluents, 625

Leaching
deltamethrin applied to cotton fabric for tsetse fly control, factors affecting its persistence, 677
PCDDs/PCDFs in flyash from incinerators to water system, 1071
polycyclic aromatic hydrocarbons from waste by solid phase extraction, 713

Mechnisms
dechlorination of chloroaromatic compds, 959
polychloroinated aromatic compds, formation with copper chloride, 987

Membranes
hexane–filled dialysis–, uptake of organochlorines from aquatic ecosystems, 997

Metabolism
2,5–dimethylhexane in male Fischer 344 rats, 77

Metals, non–metals and compounds
butyltins in estuarine sediments two years after tributyltin use was restricted, 1113
Ca ionophore and thrombin induced activation, rabbit platelets,inhibition by rubber additives, 859
Cu, Na, K and Zn in solid waste flyash from incinerators, effect on *de novo* synthesis of PCDD, 473
Cu toxicity and distribution in aquatic microcosm under different alkalinity and hardness, 577
metal contaminants. concentration USA great lakes, 809
Ni mediated biochemical alterations alleviation by cyclic and linear tetramines, 272
quantitative measurement and identification of organic constituents in hazardous waste dumps, 913
sodium methyl siliconate, photodechlorination of PCBs, 189
titanium dioxide aqueous suspension, steady–state concentration of OH radical using n–butyl chloride as probe, 1003
ZnO, irradiation with aqueous soln of 2,4–dichlorophenol, chloroaromatic products analysed by HPLC and GC–MS, 1053

Microorganisms
Pseudomonas and pure strain of *Pseudomonas cepacia*, aerobic cultures and batch cultures of PCBs, biodegradation, 3
Rhizobium, *Pisum sativum* inoculated with different strains, effect of combined nitrogen on dinitrogen fixation and productivity, 1153

Microscopy
polarised light–, and dispersion staining, asbestos in sewage sludes from 51 cities in USA, 963

Models
calculations and measure–ments, uptake and translocation of carbamates by bean plants, 327
evaluation of propanil as potential hazard to ground water, 637
for estimating prior exposure of Vietnam veterans to herbicide, 2,3,7,8–TCDD excretion, 597

phenomenological–, chlorinated hydrocarbon emission from modular incinerators, 121
predictive QSAR–, evaluating side effects of phenylureas on mammals, 613
radiative convective climate–, one dimensional time dependent, theory and development, 383
rate of transformation, pentachlorophenol in troposphere by sunlight and hydroxyl radical attack, 305
sediments in aquatic systems and clay minerals, sorption of HEDP and NTA, 15
Thermaikos gulf, Greece, runoff losses of herbicides and pesticides, 645

Molecular connectivity
use in prediction of aqueous solubility and octanol–water partition coeff, carcinogens, 723

Mutagenicity
rubber additives and inhibition of Ca ionophore and thrombin induced activation of rabbit platelets, 859

Mutagens
3–chloro–4–(dichloromethy l)–5–hydroxy–2(5H)–fur anone and related compds, formation via chlorination of phenols, 547
ether insoluble organic compds in raw and treated water extracts, 1089

Nitrates
ions, excitation induced phototransformation of monochlorophenols to chlorodihydroxybenzenes, 1063

Nitro compounds
nitrophenols and nitrobenzene, aqueous, degradation by Fentons reaction, 529

Nitrogen fixation
dinitrogen fixation and productivity in *Pisum sativum* inoculated with different strains of *Rhizobium*, effect of combined nitrogen, 1153

Oils
cooking–, biological effects of polychlorinated compds in Yusho diseaase, 537
lubricating–, in hazardous wasye dumps, 913
motor oil, fats and sediments, chromatographic automated detn of furans and dioxins, 437
motor oils, sediments and fats, retention of PCFs, PCDs and PCBs, 421
products, pyrolysis of polyethyiene–polystyrene mixtures, mass spectrometry and multidimensional chromatography, 887

Organic compounds
compds, reaction with montmorillonite, adsorptive and catalytic properties, 769
ether insoluble–, in raw and treated water extracts, genotoxic activity, 1089
hydrophobic–, partition coeff in 1–octanol/water, critical evaluation of measurement, 1175
hydrophobic–. solubility in partially miscible solvents, 939
identification of constituents in hazardous waste dumps, 913
polybrominated chemical residues, in dolphins from 1987/1988 mass mortality, US Atlantc coast, 1083
polycyclic–. bioassay of flyash using cytochrome P450IA1 induction, mouse hepatoma cell culture, 895

Ozone
increasing tropospheric concentrations, global consequences, 685

Paints
butyltins in estuarine sediments two years after tributyltin use was restricted, 1113

Partition coefficients
1–octanol/water–, measurement with hydrophobic compds, critical evaluation, 1175
n–octanol/water–, aqueous solubility estimated by bootstrap method, 953
octanol–water–, and aqueous solubility, correlation based on molecular structure, 723

Partitioning
water soluble azo dyes in activated sludge process, 107

Permeation
and sorption to phenylurea herbicides of cuticles of fruit and leaves. effects of cuticular characteristics and climatic parameters, 85

Peroxides
di–t–butyl peroxide presence, mechanism by thermolysis, acetone–induced photode–chlorination of aroclor 1254 in alkaline 2–propanol, 1161

Pesticides
air pollution in Arctic by organochlorine compds, 165
atrazine non–extractable (bound) residues in soil size fractions, localisation, 1131
degradation in soils in presence of clays or organic colloids, 769
dicofol–, urinary excreation of 4,4'–dichlorobenzilic acid by applicators, 365

ethophos, formation of interlayer compds with grafitic oxide, FTIR and gravimetric adsorption analysis, 905
inorganic and organic contaminants. concentration USA great lakes, 809
phenylureas, predictive QSAR model for evaluating side effects on mammals, 613
propanil, evaluation of potential hazard to ground water, 637
runoff losses into Thermaikos gulf, Greece, 645
surface residue from indoor fogger use, dermal transfer measured by CDFA method, 975
toxicity, from river water to aquatic organisms using *Paratya compressa improvia*, 229
uptake and translocation of–, by bean plants, model calculations and measurements, 327

Petroleum
kerosene treated rats, enzyme activity in lungs, 705
tar residues in hazardous waste dumps, 913

Phenols
2,4–dichlorophenol, irradiation of aqueous soln with ZnO, chloroaromatic products analysed by HPLC and GC–MS, 1053
chlorination, formation of 3–chloro–4–(dichloromethyl)–5–hydroxy–2(5H)–furanone and related compds, 547
halogenated anisoles in air, biohalogenation as source, 557
monochlorophenols, phototransformation to chlorodihydroxybenzene induced by excitation of nitrate ions, 1063
nitrophenols and nitrobenzene, aqueous, degradation by Fentons reaction, 529
pentachlorophenol containing solids and liquids, pyrolysis and combustion, formation of PCDDs/PCDFs, 1029
pentachlorophenol, extraction from aqueous and soil systems, comparison of methods used, 99
pentachlorophenol, transformation rate in troposphere by sunlight and hydroxyl radical attack, 305
polyhydroxybenzenes, cleavage with nitrous acid, formation of cyanide ion or cyanogen chloride, 267

Photochemistry
dechlorination of chlorobiphenyls by sodium methyl siliconate, 189
degradation, brominated dibenzodioxins and furans in organic solvents, 821
pentachlorophenol, sunlight and hydroxyl radical transformation rate in troposphere, 305
singlet oxygen in atmosphere, detn method by use of α–terpinene, 1211
toxicity of α–terthienyl towards fish in lab decreases with time, 699
transformation of monochlorophenols to chlorodihydroxybenzenes induced by excitation of nitrate ions, 1063
use of visible light in photoreduction of chloroaromatic compds, 959

Plants
bean–, uptake and translocation of carbamates, model calculations and measurements, 327
cuticles of fruit and leaves. permeation and sorption of phenylurea herbicides, effects of cuticular characteristics and climatic parameters, 85
individual response and allometry, examination of field data for sublethal toxicological effects, 747
Pisum sativum inoculated with different strains of *Rhizobium*, effect of combined nitrogen on dinitrogen fixation and productivity, 1153

Plasticisers and Plastics
##a–terthienyl extracted from soln into plastic of vessels, 699

Pollutants
aquatic–, monitoring using fish bile and radioactively labelled 4,5,6–trichloroguaiacol, 39
inorganic and organic contaminants. concentration USA great lakes, 809
micro–, 'de novo' synthesis in pilot incinerator, 1045

Pollution
air–, detn method of singlet oxygen by use of α–terpinene, 1211
air–, in Arctic by organochlorine compds, 165
atmospheric–, by auxin–type herbicides in Tala valley, Natal, 137
prevention, phenomenological models of chlorinated hydrocarbon emission from modular incinerators, 121

Polymers
polyethylene–polystyrene mixtures, pyrolysis, mass spectrometry and multidimensional chromatography, oil products, 887

Polystyrenes
mechanism of formation of polychlorinated aromatic compds with copper chloride, 987

polyethylene, part per quadrillion levels of 2,3,7,8–TCDF and 2,3,7,8–TCDD, procedure and initial results, 215

Pyrolysis
benezene–naphthalene mixtures, 279
pentachlorophenol containing solids and liquids, formation of PCDDs/PCDFs, 1029
polyethylene–polystyrene mixtures, mass spectrometry and multidimensional chromatography of oil products, 887

QSARs see Structure activity relationship

Radicals
OH radical, steady–state concentration in titanium dioxide aqueous suspension using n–butyl chloride as probe, 1003
sunlight and hydroxyl radical, rate of transformation of pentachlorophenol, 305

Reagents
Fentons reaction, degradation of aqueous nitrophenols and nitrobenzene, 529

Reduction
PCDD concentration on contaminated surfaces of PCB capacitator, 485
PCDD, PCDF emissions from waste incinerator by full scale denox unit, 569
photo–, chloroaromatic compds, use of visible light, 959

Retention indices
by gas chromatography for bromo– and bromo–chloro–dioxins, 455

Risk assessment
phase distribution and particle size of polycyclic aromatic hydrocarbons in urban atmosphere, 503

Rubber
additives, inhibition of Ca ionophore and thrombin induced activation of rabbit platelets, 859

Sediments
catalytic properties of montmorillonite in petroleum genesis, 769
chromatographic automated detn of furans and dioxins, 437
estuarine–, butyltins two years after tributyltin use was restricted, 1113
in aquatic model systems and clay minerals, sorption of HEDP and NTA, 15
inorganic and organic contaminants. concentration, USA great lakes, 809
retention of PCFs, PCDs and PCBs, 421
STORET data, toxic chemical deposition levels, regional variation, USA, 835
toxicity of *D.magna* and *C.Tentans*, effect of storage time, 801

Sewage
asbestos in sludges from 51 cities in USA, 963

Sludge
activated process–, fate of water soluble azo dyes, 107
activated–, 2,3,5–trichloro– and 2,3,6–trichloro–p–cymenene in spent bleach liquor, FTIR and mass spectral data, 881

Soil
and aqueous systems, extraction of penta–chlorophenol, comparison of methods used, 99
and fly ash, concentration of polycyclic aromatic hydrocarbons, 713
PCDD/PCDF contaminated area, induction of hepatic P450–dependent mono–oxygenase in feral mice, 1011
size fractions, localisation of atrazine non–extractable (bound) residues, 1131
sorption of nonpolar and polar compds, applicability of modified OECD–guideline, processes, measurements and experience, 106

Solubility
aqueous–, and octanol–water partition coeff, correlation based on molecular structure, 723
aqueous–, estimation from n–octanol/water partition coeffs by bootstrap method, 953
hydrophobic compds in partially miscible solvents, 939

Solvents
organic–, photodegradation of brominated dibenzodioxins and furans, 821
partially miscible–, solubility of hydrophobic compds, 939

Sorbents
evaluation, collection and analysis, trace levels of airborne bis(2–chloro ethyl)sulfide, 461

Sorption
and permeation to phenylurea herbicides of cuticles of fruit and leaves. effects of cuticular characteristics and climatic parameters, 85
chelating agents (HEDP and NTA) onto clay minerals and sediments in aquatic model systems, 15
chlorinated–benzene to hydrophobic surfaces, temp effects, 1141

nonequilibrium–, organic chemicals by low organic–carbon aquifer materials, 341
nonpolar and polar compds to soils, applicability of modified OECD–guideline, processes, measurements and experience, 106

Spectrometry
and multidimensional chromatography, analysis oil products, pyrolysis of polyethylene–polystyrene mixtures, 887
chloroaromatic products formed by irradiation of ZnO with aqueous soln of 2,4–dichlorophenol, 1053
isotope dilution HRGC/HRMS, coplanar PCBs in fish and marine animal tissue, 849
mass and FTIR data, 2,3,5–trichloro–and 2,3,6–trichloro–p–cymene ne in spent bleach liquor from activated sludge, 881
mass, composition of polycyclic aromatic hydrocarbon mixtures in uncontrolled hazardous waste dumps, 317
mass–, coplanar PCB residues, human adipose tissue from Ontario municipalities, 1019
mass–, presence of chlorinating species in turbulent flame combustion of methane with HCl, 67
quantitative measurement and identification of organic constituents in hazardous waste dumps, 913

Spectrophotometry
quantitative measurement and identification of organic constituents in hazardous waste dumps, 913

Steroids
dermal transfer of surface pesticide residue by use of indoor fogger measured by CDFA method, 975

Structure–Acivity relationship
montmorillonite, reaction with organic compds, adsorptive and catalytic properties, 769
predictive QSAR model, evaluating side effects of phenylureas on mammals, 613
QSAR models, reliability tested by bootstrap method, aqueous solubility estimated from n–octanol/water partition coeff, 953

Surfactants
anionic–, comparative biodegradability in synthetic and natural test systems, 873

Symbiosis
Pisum sativum–, effect of nitrate tolerance by *Rhizobium* on growth, nodulation and N and P content, 1153

Techniques
CDFA method, dermal transfer of surface pesticide residue by use of indoor fogger, 975
speciation and quantitation of Aroclors in hazardous wastes based on PCB congener data, 147

Terpenes
α–terpinene, use in detn method of singlet oxygen in atmosphere, 1211
chloroterpenes and other organochlorines in Baltic, Finnish and Arctic wildlife, 47
toxaphene in fish from Northern Europe, 57

Tests
leachability of polycyclic aromatic hydrocarbons from waste by solid phase extraction, 713
synthetic and natural systems, comparative biodegradability of anionic surfactants, 873

Thermal reactions
effect of mobile phase and temp on retention of dioxins, furans and PCBs in environmental and biological samples, 421
formation of halogenated hydrocarbons from methane, HCl and oxygen, 929
gasification of sorbed wastes, carbon re–activation by chemchar process, 1191

Tissues and Tissue fluids
fish and marine animal–, isotope dilution HRGC/HRMS, coplanar PCBs, 849
human adipose tissue from Ontario municipalities, coplanar PCB residues, 1019
human lymphoblastoid cells, fluorescein formation and drug metabolism, relationship to aryl hydrocarbon hydroxylase activity and environmental factors, 739
lungs, enzyme activity in kerosene treated rats, 705
plasma from several tissues of acidotic rats, effect on phosphate dependent glutaminase activity, 521
rabbit platelets, Ca ionophore and thrombin induced activation inhibition by rubber additives, 859

Toxicity
α–terthienyl towards fish in near UV–light in lab decreases with time, 699

acute–, and food avoidance behaviour of mallards and bobwhites exposed to compd 1080, 655
and bioconcentration of volatile aromatic hydrocarbons, relationship between by *Selenastrum capricornutum*, 665
and distribution of Cu in aquatic microcosm under different alkalinity and hardness, 577
chemical residues, in dolphins from 1987/1988 mass mortality, US Atlantc coast, 1083
ether insoluble organic compds in raw and treated water extracts, 1089
fluoride ion, in soft water, short–term effect to rainbow and brown trout, 603
nickel mediated biochemical changes, reversal by cyclic and linear tetramines, 272
pesticide–, from river water to aquatic organisms using *Paratya compressa improvia*, 229

Troposphere
increasing ozone concentrations, global consequences, 685
rate of sunlight and hydroxyl radical transformation of pentachlorophenol, 305

Ureas
phenylureas, predictive QSAR model for evaluating side effects on mammals, 613

Vapour and vapourisation
trace levels of airborne bis(2–chloroethyl sulfide, sorbents for analysis and collection, 461

Warfare
bis(2–chloroethylsulfide, airborne, sorbents for analysis and collection, 461

Wastes
hazardous dumps, quantitative measurement and identification, organic constituents, 913
hazardous–, speciation and quantitation of Aroclors in based on PCB congener data, 147
medical–, phenomeno–logical models of chlorinated hydrocarbon emission from modular incinerators, 121
metal contaminated–, leachability of polycyclic aromatic hydrocarbons by solid phase extraction, 713
solid waste incineration, effect of Cu, Na, K and Zn in flyash on *de novo* synthesis of PCDD, 473
sorbed–, thermal gasification, carbon re–activation by chemchar process, 1191
uncontrolled hazardous dumps, composition of polycyclic aromatic hydrocarbon mixtures, 31 7

Water
aquatic microcosm with different alkalinity and hardness, toxicity and alkalinity of Cu, 577
ground water, evaluation of propanil as potential hazard, 637
ground water, STORET data, toxic deposition, regional variation, USA, 835
humic and drinking–, formation of 3–chloro–4–(dichloromethyl)–5–hydro xy–2(5H)–furanone and related compds, via chlorination of phenols, 547
raw and treated extracts, genotoxic activity of ether insoluble organic compds, 1089
water system, PCDDs/PCDFs in flyash leached from incinerators, 1071

AUTHOR INDEX VOL. 22, 1991

Aalbers Th.G. see Bauw D.H. 713

Adams J.R. see Walling J.F. 1103

Addink R. and **Olie K.** Corrigendum 799

Albanese R.A. The chemical 2,3,7,8–tetrachlorodibenzo-*p*–dioxin and U.S. army Vietnam–era veterans 597

Albanis T.A. Runoff losses of EPTC, Molinate, Simazine, Diuron, Propanil and Metolachlor in Thermaikos Gulf, N. Greece 645

Aldrich J.C. The use of allometry and individuality in the exam– ination of field data for sublethal toxicological effects 747

Alleyne G.A.O. see Hortelano P. 521

Altschuh J. see Halfon E. 953

Anderson D. see Othoudt R.A. 801

Anderson D. see Verbrugge D.A. 809

Andreux F. see Barriuso E. 1131

Arif J.M. Khan S.G. Aslam M. Mahmood N. Joshi L.D. and **Rahman Q.** Early biochemical changes in kerosene exposed rat lungs 705

Arnoldsson see Wachtmeister C.A. 39

Asano K. see Ogawa S. 1211

Aslam M. see Arif J.M. 705

Babut M. see Vasseur P. 625

Bacci E. see Gaggi C. 869

Baek S.O. Goldstone M.E. Kirk P.W.W. Lester J.N. and **Perry R.** Phase distribution and particle size dependency of polycyclic aromatic hydrocarbons in the urban atmosphere 503

Bales R.C. see Szecsody J.E. 1141

Ballschmiter K. see Walter B. 557

Banares–Munoz M.A. see Sanchez Escribano V. 905

Barriuso E. Schiavon M. Andreux F. and **Portal J.M.** Localization of atrazine non–extractable (bound) residues in soil size fractions 1131

Bauw D.H. de Wilde P.G.M. Rood G.A. and **Aalbers Th.G.** A standard leaching test, including solid phase extraction, for the determination of PAH leach ability from waste materials 713

Behari J.R. see Srivastava S.K. 273

Belton T. see Rappe C. 239

Benfenati E. Mariani G. Fanelli R. and **Zuccotti S.** "De Novo" synthesis of PCDD, PCDF, PCDF,PCB,PCN and PAH in a pilot incinerator 1045

Bergqvist P.–A. see Rappe C. 239

Bicking M.K.L. and **Wilson R.L.** High performace size exclusion chromatography in environmental method development—1. The effect of mobile phase and temperature on the retention of dioxins, furans, and poly– chlorinated biphenyls 421

Bicking M.K.L. and **Wilson R.L.** High performce size exclusion chromatography in environmental method development—11. An automated procedure for determination of dioxins and furans 437

Bidleman T.F. see Cotham W.E.,Jr. 165

Birchard G.F. see Phillips L.J. 835

Bock K.W. see Schrenk D. 1011

Bombick D.D. see Serve M.P. 77

Boos R. Scheidl K. Prey T. Wurst F. and **Kuna R.P.** Reduction of PCDD/PCDF emissions from a waste incineration facility by a full scale DeNox–unit 569

Borwitzky H. and **Schramm K.–W.** Reduction of dioxin concentrations on contaminated surfaces 485

Boule P. see Schedel G. 1063

Boule P. see Sehili T. 1053

Brennan J.F. see Wuertz S. 1113

Brunner H. see Schrenk D. 1011

Brusseau M.L. and **Reid M.E.** Nonequilibrium sorption of organic chemicals by low organic–carbon aquifer materials 341

Bruggemann R. see Halfon E. 953

Bunce N.J. Nakai J.S. and **Yawching M.** A model for estimating the rate of chemical transformation of a VOC in the troposphere by two pathways: photolysis by sunlight and hydroxyl radical attack 305

Butterworth B.C see Kuehl D.W.

Caba J.M. see Herves A. 1153

Calxach J. see Romero J. 1089

Carson D.B. see Gledhill W.E. 873

Chamel A. see Chaumat E. 85

Chaumat E. and Chamel A. Sorption and permeation to phenylurea herbicides of isolated cuticles of fruit and leaves. Effects of cuticular characteristics and climatic parameters 85

Chemello G. see Gaggi C. 869

Chessells M. Hawker D.W. and Connell D.W. Critical evaluation of the measurement of the 1– octanol/water partition coefficient of hydrophobic compounds 1175

Cocke D.L. Ortego J.D. and Kowalska M. Interactions of montmorillonite with organic compounds – adsorptive and catalytic properties

Collins B. see Turle R. 201

Connell D.W. see Chessells M. 1175

Cooney J.J. see Wuertz S. 1113

Cotham W.E.,Jr. and Bidleman T.F. Estimating the atmospheric deposition of organochlorine contaminants to the Arctic 165

Cramm R.H. see Nestrick T.J. 215

Cristalli A. see Perez G. 279

Daughtrey E.H. see Walling J.F. 1103

de Beer P.R. see Sandmann E.R.I.C. 137

de Wilde P.G.M. see Bauw D.H. 713

Del Re A.A.M. see Trevisan M. 637

Demeter A. see Hawari J. 1161

Demeter A. see Hawari J. 189

Dempsey C.R. see Shaul G.M. 107

Deshmukh S.N. see Nigg H.N. 365

Donnelly J.R. and Sovocool G.W. Predictions of bromo– and bromochloro–dioxin GC elution properties 455

Doolittle M.M. see Wuertz S. 1113

Dostal K.A. see Shaul G.M. 107

Drake J.B. see Othoudt R.A. 801

Drake J.B. see Verbrugge D.A. 809

Draper W.M. Wijekoon D. and Stephens R.D. Speciation and quantitation of Aroclors in hazardous wastes based on PCB congener data 147

Eadon G.A. see Narang A.S. 1029

Epling G.A. Wang Q. and Qiu Q. Efficient utilization of visible light in the photoreduction of chloroaromatic compounds 959

Fanelli R. see Benfenati E. 1045

Fava F. Zappoli S. Marchetti L. and Morselli L. Biodegradation of chlorinated biphenyls (Fenclor 42) in batch cultures with mixed and pure aerobic cultures 3

Ferard J.F. see Vasseur P. 625

Fischer K. Sorption of chelating agents (HEDP and NTA) onto mineral phases and sediments in aquatic model systems–I. Sorption onto clay minerals 15

Fishman J. The global consequences of increasing tropospheric ozone concentrations 685

Fong H.R. see Ross J. 975

Forlin L. see Wachtmeister C.A. 39

Frommberger R. Polychlorinated dibenzo-*p*-dioxines and poly–chlorinated dibenzofurans in fish from South–West Germany: River Rhine and Neckar 29

Fujita S. see Ogawa S. 1211

Fukui S. see Ogawa S. 1211

Gaggi C. Chemello G. and Bacci E. Mercury vapour accumulation in azalea leaves 869

Garcia–Salguero L. see Hortelano P. 521

Ghebbioni C. see Trevisan M. 637
Giesy J.P. see Othoudt R.A. 801

Giesy J.P. see Verbrugge D.A. 809

Gledhill W.E. Trehy M.L. and Carson D.B. Comparative biodegradability of anionic surfactants in synthetic and natural test systems 873

Goldstone M.E. see Back S.O. 503

Goto S. see Mori Y. 495

Green A.E.S. Wagner J.C. and Lin K.J. Phenomenological models of chlorinated hydrocarbons 121

Greer C. see Hawari J. 1161

Grimalt J.O. see Navarro A. 913

Grimalt J.O. see Villanueva J. 317

Grzyb K.R. see Othoudt R.A. 801

Grzyb K.R. see Verbrugge D.A. 809

Haebler R. see Kuehl D.W. 1071

Hagenmaier H. see Schrenk D. 1011

Halfon E. Altschuh J. Bruggemann R. and Karcher W. Estimation of aqueous solubility from N–octanol/water partition coefficients analyzed by the bootstrap method 953
Hanasaki Y. see Ogawa S. 1211

Hatakeyama S. Shiraishi H. and Sugaya Y. Monitoring of overall pesticide toxicity of river water to aquatic organisms using a freshwater shrimp, *Paratya compressa improvisa* 229

Hawari J. Tronczynski J. Demeter A. Samson R. and Mourato D. Photodechlorination of chlorobiphenyls by sodium methyl siliconate 189

Hawari J. Demeter A. Greer C. and Samson R. Acetone–induced photodechlorination of Aroclor 1254 in alkaline 2–propanol: probing the mechanism by thermolysis in the presence of di–t– butyl peroxide 1161

Hawker D.W. see Chessells M. 1175

Heinonen P. see Herve S. 997

Herman D.C. Mayfield C.I. and Inniss W.E. The relationship between toxicity and bioconcen– tration of volatile aromatic hydrocarbons by the alga *Selenastrum capricornutum* 665

Herve S. Paukku R. Paasivirta J. Heinonen P. and Sodergren A. Uptake of organochlorines from lake water by hexane–filled dialysis membranes and by mussels

Herves A. Caba J.M. Ligero F. and Lluch C. Effect of combined nitrogen on dinitrogen fixation and productivity in *Pisum sativum* L. inoculated with different strains of *Rhizobium* 1153

Hinton W.S. and Lane A.M. Characteristics of municipal solid waste incinerator fly ash promoting the formation of polychlorinated dioxins 473

Hirohata T. Kuratsune M. Kiyohara C. and Nagayama J. Interrelationship of cellular fluorescein formation ability, an indication of cell viability, with environmental factors and with aryl hydrocarbon hydroxylase 739

Hochstein J.R. see Kononen D.W. 655

Hoke R.A. see Othoudt R.A. 801

Hoke R.A. see Verbrugge D.A. 809

Holdsworth T.J. see Shaul G.M. 107

Holmbom B. see Langvik V.–A. 547

Hormi O. see Langvik V.–A. 547

Hortelano P. Garcia–Salguero L. Alleyne G.A.O. and LupianezJ.A. Comparative effects of plasma from acutely acidotic rats on the activity of phosphate–dependent gluta– minases from several animal tissues 521

Hussain M. and Perschke H. A study of factors affecting the persistence of deltamethrin applied to cotton fabric for tsetse fly control 677

Hutzinger O. see Lenoir D. 821

Inniss W.E. see Herman D.C. 665

Inoi M. see Yamazaki H. 859

Jenkins E.J. see Manos C.G. 963

Jin H. Zhang Y. and Yang R. Toxicity and distribution of copper in an aquatic microcosm under different alkalinity and hardness 577

Joshi L.D. see Arif J.M. 705

Jyotsna see Srivastava S.K. 273

Kagan J. Stephens N. Kulakkattolickal A.T. and Wang T.P. Variables affecting the phototoxicity of a–tethienyl toward fish in the laboratory 699

Kahn P.C. see Rappe C. 239

Kakiuchi Y. see Yamazaki H. 859

Kallman B. see Pedersen J.R. 67

Kalsch W. Nagel R. and Urich K. Uptake, elimination, and bioconcentration of ten anilines in zebrafish (*Brachydanio rerio*) 351

Kanno S. see Ohya T. 267

Karcher W. see Halfon E. 953

Karenlampi S. see Kopponen P. 895

Kashimoto T. see Takaymama K. 537

Katiyar S.S. see Srivastava S.K. 273

Kawaguchi H. Steady–state concentrations of hydroxyl radical in titanium dioxide aqueous suspensions 1003

Kettrup A. see Zier B. 1121

Khalil M.A.K. see MacKay R.M. 383

Khan S.G. see Arif J.M. 705

Kim S.–W. see Koo J.–K. 887

Kinner L.L. see McGowin A.E. 1191

Kirk P.W.W. see Back S.O. 503

Kiyohara C. see Hirohata T. 739

Kjeller L.–O. see Rappe C. 239

Klein W. see von Oepen B. 285

Kononen D.W. Hochstein J.R. and Ringer R.K. Mallard and northern bobwhite exposure to Compound 1080: acute toxicity and food avoidance behavior 655

Koo J.–K. Seo Y.–H. and Kim S.–W. Multidimensional chromatography/mass spec– trometry: a tool for analysis of oil products from pyrolysis of polyethylene–polystyrene mixtures 887

Kopponen P. Torronen R. Tarhanen J. Ruuskanen J. and Karenlampi S. Cytochrome P450IA1 induction in mouse hepatome cell culture as an indicator of polycyclic organic compounds in fly ash 895

Kordel W. see von Oepen B. 285

Kowalska M. see Cocke D.L. 769

Krieger R. see Ross J. 975

Kuehl D.W. Haebler R. and Potter C. Chemical residues in dolphins from the U.S. Atlantic Coast including Atlantic bottlenose obtained during the 1987/88 mass mortality 1071

Kuehl D.W. Butterworth B.C. Libal J. and Marquis P. An isotope dilution high resolution gas chromatographic–high resolution mass spectrometric method for the determination of coplanar polychlorinated biphenyls: application to fish and marine mammals 849

Kulakkattolickal A.T. see Kagan J. 699

Kuna R.P. see Boos R. 569

Kuokkanen T. Model compounds of spent bleach liquor: 2,3,6–trichloro–p–cymenene and 2,3,5–trichloro–p–cymenene 881

Kuratsune M. see Hirohata T. 739

Lahaniatis E.S. see Zier B. 1121

Lamparski L.L. see Nestrick T.J. 215

Lane A.M. see Hinton W.S. 473

Langvik V.–A. Hormi O. Tikkanen L. and Holmbom B. Formation of the mutagen 3–chloro–4–(dichloromethyl)–5–hydroxy –2(5H)–fur anone and related compounds by chlori– nation of phenolic compounds 547

Larsen D.W. see McGowin A.E. 1191

Larsson J. see Wachtmeister C.A. 39

LeBel G.L. see Williams D.R. 1019

Lee L.S. see Pinal R. 939

Lemaire J. see Sehili T. 1053

Lenoir D. Schramm K.–W. Hutzinger O. and **Schedel G.** Photochemical degradation of brominated dibenzo–p–dioxins and –furans in organic solvents 821

Lenoir D. see Schedel G. 1063

Lenoir D. see Zier B. 1121

Lester J.N. see Baek S.O. 503

Libal J. see Kuehl D.W. 849

Ligero F. see Herves A. 1153

Lilla E. see Perez G. 279

Lin K.J. see Green A.E.S. 121
Lipczynska–Kochany E. Degradation of aqueous nitro– phenols and nitrobenzene by means of the fenton reaction 529

Lipp H.–P. see Schrenk D. 1011

Lisk D.J. see Manos C.G. 963

Liuch C. see Herves A. 1153

Lockwood K. see Rappe C. 239

Lupianez J.A. see Hortelano P. 521

MacKay R.M. and **Khalil M.A.K.** Theory and development of a one dimensional time dependent radiative convective climate model 383

Mahmood N. see Arif J.M. 705

Manahan S.E. see McGowin A.E. 1191

Manos C.G. Patel–Mandlik K.J. Ross B.J. Jenkins E.J. and **Lisk D.J.** Prevalence of asbestos in sewage sludges from 51 large and small cities in the United States 963

Marchetti L. see Fava F. 3

Margetich S. see Ross J. 975

Mariani G. see Benfenati E. 1045

Marquis P. see Kuehl D.W. 849

Matsushita H. see Mori Y. 495

Mattle D.R. see Serve M.P. 77

Mayfield C.I. see Herman D.C. 665
McDonald G.A. see Serve M.P. 77

McGowin A.E. Kinner L.L. Manahan S.E. and **Larsen D.W.** ChemChar process for carbon re–activation 1191
Miller M.E. see Wuertz S. 1113
Mimura M. see Takayama K. 537

Miyata H. see Takayama K. 537

Montepiani C. see Trevisan M. 637

Mori Y. Goto S. Onodera S. Naito S. and **Matsushita H.** Aqueous chlorination of tetracyclic aromatic hydro– carbons: reactivity and product distribution 495

Morselli L. see Fava F. 3

Mourato D. see Hawari J. 189

Nagayama J. see Hirohata T. 739

Nagel R. see Kalsch W. 351

Naito S. see Mori Y. 495

Nakai J.S. see Bunce N.J. 305

Narang A.S. Swami K. Narang R.S. and **Eadon G.A.** Pyrolysis and combustion of liquids and solids containing pentachloro– phenol 1029

Narang R.S. see Narang A.S. 1029

Navarro A. Rosell A. Villanueva J. and **Grimalt J.O.** Monitoring of hazardous waste dumps by the study of metals and solvent– soluble organic chemicals 913

Navarro A. see Villanueva J. 317

Nendza M. Predictive QSAR models estimating ecotoxic hazard of phenylureas: mammalian toxicity 613

Nestrick T.J. Lamparski L.L. and **Cramm R.H.** Procedure and initial results for the deter– mination of low part per quadrillion levels of 2,3,7,8–tetrachlorodibenzo–p–dioxin and 2,3,7,8–tetrachlorodibenzofuran in polyethylene 215

Nigg H.N. Stamper J.H. Deshmukh S.N. and **Queen R.M.** 4,4'–Dichlorobenzilic acid urinary excretion by dicofol pesticide applicators 365

Norstrom R.J. see Turle R. 201

Ogawa S. Fukui S. Hanasaki Y. Asano K. Uegaki H. Fujita S. and **Shimazaki R.** Determination method of singlet oxygen in the atmosphere by use of a–terpinene 1211
Ohta S. see Takayama K. 537

Ohya T. and **Kanno S.** Formation of cyanide ion or cyanogen chloride through the cleavage of aromatic rings by nitrous acid or chlorine–XII. On the reactions of polyhydroxybenzenes with nitrous acid 267

Olie K. see Addink R. 799

Olsson J.O. see Pedersen J.R. 67

Onodera S. see Mori Y. 495

Ortego J.D. see Cocke D.L. 769

Othoudt R.A. Giesy J.P. Grzyb K.R. Verbugge D.A. Hoke R.A. Drake J.B. and **Anderson D.** Evaluation of the effects of storage time on the toxicity of sediments 801

Othoudt R.A. see Verbrugge D.A. 809

Paasivirta J. and **Rantio T.** Chloroterpenes and other organochlorines in Baltic, Finnish and Arctic wildlife 47

Paasivirta J. see Herve S. 997

Patel–Mandlik K.J. see Manos C.G. 963

Patil G.S. Correlation of aqueous solubility and octanol–water partition coefficient based on molecular structure 723

Paukku R. see Herve S. 997

Pedersen J.R. Kallman B. Simonson M. and **Olsson J.O.** The chlorinating species in turbulent flame combustion of mathane with hydrogen chloride present 67

Pedersen J.R. A parametric study of the homogeneous reaction between methane, hydrogen chloride and oxygen 929

Perez G. Cristalli A. and **Lilla E.** Pyrolysis of benzene–naph– thalene mixtures 279

Perry R. see Baek S.O. 503

Perschke H. see Hussain M. 677

Phillips L.J. and **Birchard G.F.** Use of storet data to evaluate variations in environmental contamination by census division 835

Pinal R. Lee L.S. and **Rao P.S.C.** Prediction of the solubility of hydrophobic compounds in nonideal solvent mixtures 939

Portal J.M. see Barriuso E. 1131

Posner J.C. Evaluation of sorbents for the collection and analysis of trace levels of airborne vapors: bis(2–chloroethyl) sulfide (mustard), a case study 461

Potter C. see Kuehl D.W. 1071

Prey T. see Boos R. 569

Pussemier L. see Trapp St. 1085

Pussemier L. see Trapp St. 327

Qiu Q. see Epling G.A. 959
Queen R.M. see Nigg H.N. 365

Rahman Q. see Arif J.M. 705

Rantio T. see Paasivirta J. 47

Rao P.S.C. see Pinal R. 939

Rappe C. Bergqvist P.–A. Kjeller L.–O. Swanson S. Belton T. Ruppel B. Lockwood K. and Kahn P.C. Levels and patterns of PCDD and PCDF contamination in fish, crabs, and lobsters from Newark Bay and the New York Bight 239

Reid M.E. see Brusseau M.L. 341

Ringer R.K. see Kononen D.W. 655

Rivera J. see Romero J. 1089

Roberts J. see Serve M.P. 77

Romero J. Ventura F. Caixach J. and Rivera J. Genotoxic activity of ether insoluble organic com– pounds in raw and treated water extracts 1089

Rood G.A. see Bauw D.H. 713

Rosell A. see Navarro A. 913

Rosell A. see Villanueva J. 317

Ross B.J. see Manos C.G. 963

Ross J. Fong H.R. Thongsinthusak T. Margetich S. and Krieger R. Measuring potential dermal transfer of surface pesticide residue generated from indoor fogger use: using the CDFA roller method interim report 975

Ruppel B. see Rappe C. 239

Ruuskanen J. see Kopponen P. 895

Samson R. see Hawari J. 1161
Samson R. see Hawari J. 189

Sanchez Conejero J.M. see Sanchez Escribano V.

Sanchez Escribano V. Banares–Munoz M.A. and Sanchez Conejero J.M. Interaction and formation of ethoprophos–graphitic oxide interlayer compounds 905

Sandmann E.R.I.C. de Beer P.R. and van Dyk L.P. Atmospheric pollution by auxin–type–herbi cides in Tala Valley, Natal 137

Schedel G. Lenoir D. and Boule P. Phototransformation of monochlorophenols induced by excitation of nitrate ions 1063

Schedel G. see Lenoir D. 821

Scheidl K. see Boos R. 569

Schiavon M. see Barriuso E. 1131

Schramm K.–W. see Borwitzky H. 485

Schramm K.–W. see Lenoir D. 821

Schrenk D. Lipp H.–P. Brunner H. Wiesmuller T. Hagenmaier H. and Bock K.W. Induction of hepatic P450–dependent mono– oxygenase in feral mice from a PCDD/PCDF–contaminated area 1011

Sehili T. Boule P. and Lemaire J. Photocatalysed transformation of chloroaromatic derivatives on zinc oxide IV: 2,4–dichloro– phenol 1053

Seo Y.–H. see Koo J.–K. 887

Serve M.P. Bombick D.D. Roberts J. McDonald G.A. Mattie D.R. and Yu K.O. The metabolism of 2,5–dimethylhexane in male Fischer 344 rats 77

Shaul G.M. Holdsworth T.J. Dempsey C.R. and Dostal K.A. Fate of water soluble azo dyes in the activated sludge process 107

Shimazaki R. see Ogawa S. 1211

Shiraishi H. see Hatakeyama S. 229

Simonson M. see Pedersen J.R. 67
Sodergren A. see Herve S. 997

Sovocool G.W. see Donnelly J.R. 455

Srivastava A. see Srivastava S.K. 273

Srivastava R.C. see Srivastava S.K. 273

Srivastava S.K. Srivastava A. Katiyar S.S. Jyotsna. Behari J.R. and Srivastava R.C. Alleviation of nickel mediated biochemical alterations by cyclic and linear tetramines 273

Stamper J.H. see Nigg H.N. 365

Stephens N. see Kagan J. 699

Stephens R.D. see Draper W.M. 147

Stieglitz L. see Jay K. 987

Stratton G.W. see Wall A.J. 99

Sugaya Y. see Hatakeyama S. 229

Swami K. see Narang A.S. 1029

Swanson S. see Rappe C. 239
Szecsody J.E. and Bales R.C. Temperature effects on chlorinated–benzene sorption to hydrophobic surfaces 1141

Tai H.–H. see Yamazaki H. 859

Takahashi K. see Takenaka S. 375

Takayama K. Miyata H. Mimura M. Ohta S. and Kashimoto T. Evaluation of biological effects of polychlorinated compounds found in contaminated cooking oil responsible for the disease "Yosho" 537

Takenaka S. and Takahashi K. Enhancement of fecal excretion of polychlorinated biphenyls by the addition of rice bran fiber to the diet in rats 375

Tarazona J.V. see Camargo J.A. 605

Tarhanen J. see Kopponen P. 895

Thongsinthusak T. see Ross J. 975

Tikkanen L. see Langvik V.–A. 547

Torronen R. see Kopponen P. 895

Trapp St. and Pussemier L. Corrigendum 1085

Trapp St. and **Pussemier L.** Model calculations and measurements of uptake and translocation of carbamates by bean plants 327

Trehy M.L. see Gledhill W.E. 873

Trevisan M. Montepiani C. Ghebbioni C. and **Del Re A.A.M.** Evaluation of potential hazard of propanil to ground–water 637
Tronczynski J. see Hawari J. 189
Turle R. Norstrom R.J. and **Collins B.** Comparison of PCB quantitation methods: re–analysis of archived specimens of Herring Gull eggs from the Great Lakes 201

Uegaki H. see Ogawa S. 1211

Urich K. see Kalsch W. 351

van der Valk F. and **Wester P.G.** Determination of toxaphene in fish from Northern Europe 57 57

van Dyk L.P. see Sandmann E.R.I.C. 137

Vasseur P. Ferard J.F. and **Babut M.** The biological aspects of the regulatory control of industrial effluents in France 625

Ventura F. see Romero J. 1089

Verbrugge D.A. Othoudt R.A. Grzyb K.R. Hoke R.A. Drake J.B. Giesy J.P. and **Anderson D.** Concentrations of inorganic and organic contaminants in sedi– ments of six harbors on the North American Great Lakes 809

Verbugge D.A. see Othoudt R.A. 801

Villanueva J. Rosell A. Grimalt J.O. and **Navarro A.** Chemical characterization of polycyclic aromatic hydrocarbon mixtures in uncontrolled hazardous waste dumps 317

Villanueva J. see Navarro A. 913

von Oepen B. Kordel W. and **Klein W.** Sorption of nonpolar and polar compounds to soils: processes, measurements and experience with the applicability of the modified OECD–Guideline 106 285
Wachtmeister C.A. Forlin L. Arnoldsson and **Larsson J.** Fish bile as a tool for monitoring aquatic pollutants: studies with radioactively labelled 4,5,6–trichloroguaiacol 39

Wagner J.C. see Green A.E.S. 121

Wall A.J. and **Stratton G.W.** Comparison of methods for the extraction of pentachlorophenol from aqueous and soil systems 99

Walling J.F. Daughtrey E.H. and **Adams J.R.** Changes in poly– cyclic aromatic hydrocarbon content of atmospheric particulate monitored using fuzzified chromatograms 1103

Walter B. and **Ballschmiter K.** Biohalogenation as a source of halogenated anisoles in air 557

Wang Q. see Epling G.A. 959

Wang T.P. see Kagan J. 699

Wester P.G. see van der Valk F. 57

Wiesmuller T. see Schrenk D. 1011

Wijekoon D. see Draper W.M. 147
Williams D.T. LeBel G.L. Coplanar polychlorinated biphenyl residues in human adipose tissue samples from Ontario municipalities 1019

Wilson R.L. see Bicking M.K.L. 421

Wilson R.L. see Bicking M.K.L. 437

Wuertz S. Miller M.E. Doolittle M.M. Brennan J.F. and **Cooney J.J.** Butyltins in estuarine sediments two years after tributyltin use was restricted 1113

Wurst F. see Boos R. 569

Yamaguchi T. see Yamazaki H. 859
Yamauchi A. see Yamazaki H. 859

Yamazaki H. Inoi M. Yamaguchi T. Yamauchi A. Kakiuchi Y. and **Tai H.–H.** Rubber additives inhibit the calcium ionophore and thrombin induced activation of rabbit platelets 859

Yang R. see Jin H. 577

Yawching M. see Bunce N.J. 305

Yu K.O. see Serve M.P. 77

Zappoli S. see Fava F. 3

Zhang Y. see Jin H. 577

Zier B. Lenoir D. Lahaniatis E.S. and **Kettrup A.** Surface catalyzed halogenation– dehalogenation reactions of aromatic bromine compounds adsorbed on fly ash 1121

Zuccotti S. see Benfenati E. 1045


1.0
1.1
1.25
1.4
1.6
2.8
2.5
3.2
2.2
3.6
2.0
4.0
1.8